



## 预防与诊断服务 情况说明

在计划筛查时，了解预防性检测与诊断性检测之间的区别很重要。根据检测的种类，您的保险范围可能会有所不同。美国癌症协会发现，有确切证据表明，在 45 至 75 岁的成年人中筛查结肠直肠癌可降低结肠直肠癌死亡率。<sup>1</sup>

### 了解预防性筛查与诊断性检测的差异及其对保险范围的影响



#### 预防性筛查

对没有症状但有结肠直肠癌风险的患者群体进行检测（例如，大便隐血试验、粪便免疫化学试验、粪便 DNA 检测 [Cologuard<sup>®</sup>]、乙状结肠镜检查或结肠镜检查），例如以下群体<sup>2,3</sup>：

- 年龄 45 - 75 岁<sup>2</sup>
- 有结肠癌家族史<sup>2</sup>

许多医疗保健计划可涵盖预防性筛查，不需要自费。请咨询您保险提供商，了解更多信息。<sup>4</sup>



#### 诊断性检测

结肠镜检查用于根据某些症状诊断健康状况，例如<sup>4,5</sup>：

- 出血或大便形状不规则
- 预防性筛查结果呈阳性

您可能需要支付共付额、共负保险或自付额。<sup>4</sup>

**立即咨询您的医疗保健提供者和保险公司。**了解哪些筛查适合您、您的计划涵盖哪些服务以及您的花费可能是多少，然后安排结肠癌筛查。

如需了解更多关于结肠直肠癌和筛查的信息，请访问

[www.LoveYourColon.org](http://www.LoveYourColon.org)

参考资料：1. Wolf AM, Fontham ET, Church TR, et al. Colorectal cancer screening for average-risk adults: 2018 guideline update from the American Cancer Society. *CA Cancer J Clin.* 2018;68(4):250-281. 2. American Cancer Society Guideline for Colorectal Cancer Screening. American Cancer Society. <https://www.cancer.org/cancer/colon-rectal-cancer/detection-diagnosis-staging/acs-recommendations.html>. Accessed April 9, 2020. 3. Colorectal cancer screening tests. American Cancer Society website. <https://www.cancer.org/cancer/colon-rectal-cancer/detection-diagnosis-staging/screening-tests-used.html>. Accessed March 23, 2020. 4. Insurance coverage for colorectal cancer screening. American Cancer Society website. <https://www.cancer.org/cancer/colon-rectal-cancer/detection-diagnosis-staging/screening-coverage-laws.html>. Accessed March 23, 2020. 5. Signs and symptoms of colorectal cancer. American Cancer Society website. <https://www.cancer.org/latest-news/signs-and-symptoms-of-colon-cancer.html>. Accessed March 23, 2020.